

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

2132.101

Examiner copy

reference

#3

About Entrez

Limits: Publication Date from 1995 to 1995

Display Abstract

Show 20

Sort by

Send to

Text Version

All: 1 Review: 0 ☒

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

☐ 1: Acta Diabetol. 1995 Dec;32(4):251-6.

Related Articles, Links

Apolipoprotein AI-CIII-AIV genetic polymorphisms and coronary heart disease in type 2 diabetes mellitus.

Rigoli L, Raimondo G, Di Benedetto A, Romano G, Porcellini A, Campo S, Corica F, Riccardi G, Squadrito G, Cucinotta D.

Department of Internal Medicine, University of Messina, Italy.

The aim of this study was to verify whether or not the increased prevalence of coronary heart disease (CHD) commonly observed in patients with type 2 diabetes mellitus is related to a genetic background involving restriction fragment length polymorphisms (RFLPs) of apolipoproteins. On the basis of a case-control design, 62 type 2 diabetic patients with CHD (confirmed by clinical history and electrocardiogram) and 62 age- and sex-matched diabetic subjects without CHD were enrolled. In each of them RFLPs of the apolipoprotein CIII gene (S1 or S2 allele) and AI promoter region (A or G allele), together with fasting plasma lipids and apolipoproteins levels, were assessed. The rare S2 allele was found significantly ($P = 0.05$) more frequently in patients with CHD, and its related S1S2 genotype was associated with higher plasma levels of total cholesterol ($P = 0.01$), triglycerides ($P = 0.007$) and apo B ($P = 0.001$) than the S1S1 genotype. The A allele was more frequent ($P = 0.004$) in patients without CHD and was associated with lower plasma cholesterol ($P = 0.0001$), low-density lipoprotein (LDL)-cholesterol ($P = 0.0001$) and apo B ($P = 0.005$). The S1/A haplotype was more frequent ($P = 0.05$) in patients without CHD and was associated with the lowest plasma lipid levels. These results suggest that genetic factors, related to the apo AI-CIII-AIV gene cluster, could play a role in the development of CHD in type 2 diabetic patients, probably through modification of their plasma lipid pattern.

PMID: 8750764 [PubMed - indexed for MEDLINE]

Display Abstract



Show 20

Sort by

Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 27 2005 04:57:20



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Limits: Publication Date from 1990 to 2000

Display Abstract

Show 20

Sort by

Send to

All: 1

Review: 0

☐ 1: Diabetes. 1994 Dec;43(12):1485-9.

Related Articles, Links

ApoA-IV polymorphism associated with myocardial infarction in obese NIDDM patients. The San Luis Valley Diabetes Study.

Rewers M, Kamboh MI, Hoag S, Shetterly SM, Ferrell RE, Hamman RF.

Department of Preventive Medicine, University of Colorado, Denver 80262.

Non-insulin-dependent diabetes mellitus (NIDDM) confers myocardial infarction (MI) risk unexplained by known factors. In 356 NIDDM patients and 1,087 people with normal glucose tolerance, we investigated the association between MI risk and polymorphism at codon 360 in the apolipoprotein A-IV (apoA-IV) gene. During 1984-1992, MI was diagnosed in 84 diabetic and in 106 nondiabetic people. The risk of MI did not differ by apoA-IV phenotype in nondiabetic people; however, in NIDDM patients, those with the apoA-IV 1-2 phenotype had 2.8 (95% confidence interval: 1.4-5.6) higher MI risk than those with the 1-1 phenotype, adjusting for age, gender, ethnicity, hypertension, smoking, body mass index, fat centrality, and low-density lipoprotein and high-density lipoprotein cholesterol. The risk of MI was particularly high in obese NIDDM patients with the apoA-IV 1-2 phenotype: 5.1 (2.4-11.2) times that in obese apoA-IV 1-1 NIDDM patients and 7.7 (3.6-16.7) times that in lean nondiabetic people. The effect of apoA-IV 1-2 did not appear to be a part of the insulin-resistance syndrome nor was it dependent on diabetes duration or control. One half of the excess MI risk in the diabetic population studied was explained by the apoA-IV 1-2 phenotype. These results indicate that approximately 17% of NIDDM patients have a high MI risk apoA-IV phenotype that is particularly deleterious in obese patients.

PMID: 7958503 [PubMed - indexed for MEDLINE]

Display Abstract

Show 20

Sort by

Send to



[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 27 2005 04:57:20



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Limits: Publication Date from 1992 to 1992

Display Abstract

Show 20

Sort by

Send to

All: 1 Review: 1 ☒☐ 1: Cancer Res. 1992 May 1;52(9 Suppl):2711s-2718s.

Related Articles, Links

Considerations in bringing a cancer biomarker to clinical application.

Tockman MS, Gupta PK, Pressman NJ, Mulshine JL.Department of Environmental Health Sciences, Johns Hopkins University
School of Hygiene and Public Health, Baltimore, Maryland 21205.

Specific challenges face our application of emerging biomarkers to early lung cancer detection. These challenges might be considered frontiers to be bridged between established biomedical disciplines, requiring expertise often beyond the range of individual investigators. Cross-disciplinary research already has led to new appreciation of the mechanisms which underlie the phenotypic expression of the transformed cell and places within our grasp the tools which might lead to successful early lung cancer detection. Prior to the successful application of newly described markers, further cross-disciplinary research must (a) refine the selection of biologically appropriate markers, (b) validate such markers against acknowledged disease end points, (c) establish quantitative criteria for marker presence/absence, and (d) confirm marker predictive value in prospective population trials.

Publication Types:

- Review
- Review, Tutorial

PMID: 1563002 [PubMed - indexed for MEDLINE]

Display Abstract

Show 20

Sort by

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Jun 27 2005 04:57:20